

AD\_\_\_\_\_

Award Number: DAMD17-01-1-0444

TITLE: Aromatase Overexpression and Breast Cancer Development

PRINCIPAL INVESTIGATOR: Doctor Wei Yue

CONTRACTING ORGANIZATION: University of Virginia  
Charlottesville, VA 22904

REPORT DATE: August 2005

TYPE OF REPORT: Annual Summary

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;  
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

20060110 080

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

1. REPORT DATE 01-08-2005			2. REPORT TYPE Annual Summary		3. DATES COVERED 1 Aug 2001 – 31 Jul 2005	
4. TITLE AND SUBTITLE  Aromatase Overexpression and Breast Cancer Development					5a. CONTRACT NUMBER	
					5b. GRANT NUMBER DAMD17-01-1-0444	
					5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)  Doctor Wei Yue					5d. PROJECT NUMBER	
					5e. TASK NUMBER	
					5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  University of Virginia Charlottesville, VA 22904					8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012					10. SPONSOR/MONITOR'S ACRONYM(S)	
					11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited						
13. SUPPLEMENTARY NOTES						
14. ABSTRACT Estrogen can be metabolized to hydroxylated catechol estrogen, a genotoxic metabolite of estrogen, which causes DNA damage and tumors in animal models. In situ synthesis of estrogen in the breast through aromatase results in high tissue estrogen concentrations. We hypothesized that overexpression of aromatase in breast tissue increases tissue estradiol concentrations and consequent genotoxic metabolites, and eventually causes breast cancer. To test our hypothesis, we stably expressed aromatase cDNA in MCF-10A cells, a benign breast epithelial cell line (MCF-10Aarom). We demonstrated that MCF-10Aarom cells expressed functional aromatase using tritiated water release assay and products isolation by thin layer chromatography. MCF-10Aarom cells, incubated for 3 months with aromatase substrate, androstenedione, formed colonies in soft agar indicating the overexpression of aromatase induces cellular transformation. MCF-10Aarom cells have all enzymes required to convert estrogen to catechoestrogens and quinine. Overexpression of aromatase enhanced production of genotoxic metabolites, which could be blocked by aromatase inhibitor, letrozole.						
15. SUBJECT TERMS aromatase, non-cancerous human breast epithelial cells, stable transfection, nude mouse, estrogen metabolites, depurinating DNA adduct, hormone carcinogen						
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON	
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			USAMRMC	
			UU	9	19b. TELEPHONE NUMBER (include area code)	

## Table of Contents

Cover.....	
SF 298.....	
Table of Contents.....	
Introduction.....	2
Body.....	2
Key Research Accomplishments.....	4
Reportable Outcomes.....	4
Conclusions.....	5
References.....	5
Career Development.....	6
Appendices.....	6

## Introduction

Clinical data and results from animal studies indicate that estrogen not only promotes breast cancer growth but also plays a role in breast cancer initiation. The carcinogenic effect of estrogen is mediated by hydroxylated catechol estrogen, a genotoxic metabolite of estrogen, which causes DNA damage (1-4). In situ synthesis of estrogen in the breast through aromatase makes the major contribution to the high tissue estrogen concentrations (5). We hypothesized that overexpression of aromatase in breast tissue increases tissue estradiol concentrations and consequent genotoxic metabolites, and eventually causes breast cancer. To test our hypothesis, we have established a stable cell line of benign breast epithelial cell by expressing aromatase gene in MCF-10A cells. During this annual report period, we measured estrogen metabolites and depurinating DNA adduct in aromatase expressing MCF-10A cells (Specific Aim 2); fulfilled the in vivo tumorigenesis studies (Specific Aim 3) and carried out an assay for loss of heterozygosity (Specific Aim 3).


## Body

### I. In vitro study


#### 1. Metabolite measurement

MCF-10A<sup>arom</sup> and MCF-10A<sup>vect</sup> cells were cultured in 150 mm dishes until confluence. The cells were treated in serum free medium with vehicle, estradiol, 4-OHE<sub>2</sub>, androstenedione plus or minus letrozole for 24 h. All compounds were at the concentration of 10  $\mu$ M. Cells and media were collected and kept at -80C until analysis. Estrogen metabolites, conjugates, and depurinated DNA adducts were measured by HPLC with a 12-channel electrochemical detector and confirmed by mass spectrometry (6-8). Figure 1 illustrates the metabolic pathways of estrogen and measurable metabolites/DNA adducts.

MCF-10A cells have all enzymes that catalyze conversion of estrogen to catechoestrogens and their methylation (Fig. 2). Incubation of MCF-10A<sup>arom</sup> cells with estradiol (10  $\mu$ M) for 24 hours dramatically increased production of 4-hydroxyestrone (Fig. 2A). Incubation with aromatase substrate, androstenedione ( $\Delta^4$ A), also increased the level of 4-hydroxyestrone (Fig. 2A) indicating that  $\Delta^4$ A was aromatized to estrogen which was metabolized to 4-hydroxyestrone. A large portion of catechoestrogens were converted to methoxyestrogens,



detoxification metabolites, by catecho-*o*-methyl transferase (COMT). This is more significant when 4-OHE<sub>2</sub> was used as a precursor (Fig. 2B). Estrogen-3,4-quinones (E-3,4-Q) are the core genotoxic metabolites of estrogens. Estrogen-3,4-quinones are highly reactive and form conjugates with glutathione or bind to purine motifs on DNA as soon as they form (Fig. 1). Formation of glutathione conjugates and their metabolites reflects production of E-3,4-Q. As shown in Fig. 2C, 4-OHE<sub>2</sub>-2-NACys, a conjugate of E-3,4-quinone, was detected in MCF-10A<sup>arom</sup> cells pretreated with vehicle, E<sub>2</sub>, 4-OHE<sub>2</sub>, and Δ<sup>4</sup>A. Depurinated DNA adduct, 4-OHE<sub>2</sub>-1-N7Gua, was only detected in the cells treated with E<sub>2</sub>, 4-OHE<sub>2</sub>, and Δ<sup>4</sup>A. The level of 4-OHE<sub>2</sub>-1-N7Gua was undetectable in the cells incubated with Δ<sup>4</sup>A plus the specific aromatase inhibitor, letrozole (Fig. 2D).



## 2. Loss of heterozygosity (LOH)

Loss of heterozygosity may contribute to the initiation and progression of breast cancer. LOH has been observed in both in situ and invasive breast carcinomas on multiple chromosomal arms (9, 10) as well as in normal tissue adjacent to breast carcinomas (11). Human breast epithelial cells transformed with estradiol also showed LOH in chromosome 11 (12). To determine whether aromatase overexpression causes gene instability, LOH was assessed in MCF-10A<sup>arom</sup> cells in collaboration with Dr. Russo at Fox Chase Cancer Center, Philadelphia.

Cell lines: MCF-10A (regular MCF-10A), MCF-10A<sup>vect</sup> (transfected with empty pH $\beta$  vector) and MCF-10A<sup>arom</sup> cells.

Treatment: MCF-10A<sup>arom</sup> cells were treated for 10 months with androstenedione ( $\Delta^4$ A,  $10^{-6}$  M),  $\Delta^4$ A + letrozole ( $10^{-7}$  M), letrozole alone and estradiol ( $10^{-6}$  M). Regular MCF-10A and MCF-10A<sup>vect</sup> cells were without treatment.

Genomic DNA was extracted LOH was assayed using microsatellite polymorphism makers TP53dint and D13S893 that flank *p53* (chromosome 17) and *WT2* (chromosome 11p15.5) respectively. Both genes are tumor suppress genes.

All samples display one allele of 185 bp for the marker D13S893 and one allele of 295 bp for the marker TP53dint. It is necessary to examine more loci.

## II. In vivo study

To determine whether aromatase overexpression causes genotoxic damage and consequently mammary tumor, tumorigenesis of aromatase expressing MCF-10 cells were assessed in nude mice. Long term experiment for tumor formation and prevention was carried out during this report period.

Ovariectomized nude mice were inoculated subcutaneously with 5 million MCF-10A<sup>arom</sup> on left flank and regular MCF-10A cells on right flank. The animals were then divided into three groups, 20 mice each. One group of animals received androstenedione silastic capsule to provide aromatase precursor. One group of animals received androstenedione plus letrozole (5  $\mu$ g/day, s.c.). The third group served as a control. The mice were treated for one year. There was no palpable tumor developed from either regular MCF-10A or from MCF-10A<sup>arom</sup> cells.

## Key Research Accomplishments

- Determination of estrogen metabolites and depurinating DNA adduct in MCF-10A<sup>arom</sup> cells
- Loss of heterozygosity assay
- in vivo tumorigenesis

## Reportable Outcomes

An abstract entitled "Investigation of the carcinogenic effect of estradiol in the in vitro and in vivo models without estrogen receptors" was submitted to the International Conference of Aromatase 2004 for poster presentation in September 2004, Edinburgh, United Kingdom.

## Conclusions

MCF-10A<sup>arom</sup> cells have all enzymes that catalyze formation of catechoestrogen and methoxyestrogen. Expression of aromatase increases in situ estrogen synthesis and its genotoxic products, which can be blocked by aromatase inhibitor, letrozole. There is no LOH detected in the loci using TP53dint and D13S893 as markers. No tumor formed from MCF-10A<sup>arom</sup> cells in nude mice.

## Reference

1. Zhu BT, Bui QD, Weisz J, and Liehr J: Conversion of estrone to 2- and 4-hydroxyestrone by hamster kidney and liver microsomes: implications for the mechanism of estrogen-induced carcinogenesis. *Endocrinology* 135: 1772-1779, 1994.
2. Roy D, Weisz J, and Liehr JG: The O-methylation of 4-hydroxyestradiol is inhibited by 2-hydroxyestradiol: implications for estrogen-induced carcinogenesis. *Carcinogenesis* 11:459-462, 1990.
3. Roy D, Bernhardt A, Strobel H, Liehr JG: Catalysis of the oxidation of steroid and stilbene estrogens of estrogen quinone metabolites by the  $\eta$ -naphthoflavone-inducible cytochrome P450 1A family. *Arch. Biochem. Biophys.* 296: 450-456, 1992.
4. Cavalieri EL, Stack DE, Devanesan PD, Todorovic R, Dwivedy I, Higginbotham S, Johansson SL, Patil KD, Gross ML, Gooden JK, Ramanathan R, Cerny RL, and Rogan EG: Molecular origin of cancer: catechol estrogen-3,4-quinones as endogenous tumor initiators. *Proc. Natl. Acad. Sci. U.S.A.* 94: 10937-10942, 1997.
5. Yue W, Wang J-P, Hamilton CJ, Demers LM, and Santen RJ: In situ aromatization enhances breast tumor estradiol levels and cellular proliferation. *Cancer Res.* 58: 927-932, 1998.
6. Cavalieri, E., Frenkel, K., Liehr, J.G., Rogan, E., Roy, D. Estrogens as endogenous genotoxic agents ---- DNA adducts and mutations. *J. Natl. Cancer Inst. Monogr.* 27:75-93, 2000.
7. Cavalieri, E.L., Kumar, S., Todorovic, R., Higginbotham, S., Badawi, A.F., Rogan, E.G. Imbalance of estrogen homeostasis in kidney and liver of hamsters treated with estradiol: implications for estrogen-induced initiation of renal tumors. *Chem. Res. Toxicol.* 14:1041-1050, 2001.
8. Rogan, E.G., Badawi, A.F., Devanesan, P.D., Meza, J.L., Edney, J.A., Wert, W.W., Higginbotham, S.M., Cavalieri, E.L. Relative imbalances in estrogen metabolism and conjugation in breast tissue of women with carcinoma: potential biomarkers of susceptibility to cancer. *Carcinogenesis* 24:697-702, 2003.
9. Radford, D.M., Fair, K.L., Phillips, N.J., Ritter, JH. Steinbrueck, T. Holt, MS. Donis-Keller, H. Allelotyping of ductal carcinoma in situ of the breast: deletion of loci on 8p, 13q, 16p, 17p and 17q. *Cancer Res.* 55:3399-3405, 1995.
10. Devilee, P. Hermans, J., Eyfjord, J. Boorresen, AL. Liderau, R. Sobol, H. Borg, A. Cleton-Jansen, AM. Olah, E. Cohen, BB. Scherneck, S. Hamann, U. Peterlin, B. Caligo, M. Bignon, YJ. Maugard, C. Loss of heterozygosity at 7q31 in breast cancer: results from an international collaborative study group. The Breast Cancer Somatic Genetics Consortium. *Genes Chromosomes Cancer* 18: 193-199, 1997
11. Deng, Guoren, Lu, You, Zlotnikov, Galiana, Thor, Ann D., Smith, Helene S. Loss of heterozygosity in normal tissue adjacent to breast carcinomas. *Science* 274:2057-2059, 1996.

12. Russo J, Lareef MH, Tahin Q, Hu Y-F, Slater C, Ao X, and Russo IH: 17 $\beta$ -Estradiol is carcinogenic in human breast epithelial cells. *J. Steroid Biochem. Mol. Biol.* 80: 149-162, 2002.

## **Career Development**

The PI was promoted to Associate Professor of Research in 2002. The PI attended International Aromatase Conference as an invited speaker in Kyoto, Japan, 2002. DOD support also allowed the PI to attend the International Conference of Aromatase 2004 in Edinburgh, UK to report findings of the research project funded by DOD. During the funding period, the PI submitted an original and revised application for NIH R01 grant.

## **Appendices**

Abstract submitted to the International Conference of Aromatase 2004.



## Investigation of the carcinogenic effect of estradiol in the in vitro and in vivo models without estrogen receptors

Wei Yue, Jiping Wang, Sandra Gunselman, Ercole Cavalieri, Eleanor Rogan, and Richard Santen

Department of Internal Medicine, University of Virginia and Eppley Institute for Cancer Research, University of Nebraska Medical Center, USA

Estrogen causes cancers in animals. Substantial correlative data suggest that estrogens cause breast cancer in women. A body of emerging data suggests that estrogen is metabolized to genotoxic products that damage DNA and initiate cancer. This mechanism together with receptor-mediated stimulation of cell proliferation may act in concert to induce breast cancer. To test the genotoxic metabolite hypothesis, two models were employed: MCF-10A benign breast epithelial cells expressing aromatase were used to determine production of estrogen metabolites and their blockade by the aromatase inhibitor, letrozole. ER $\alpha$  knockout mice expressing the Wnt-1 oncogene (ERKO/Wnt-1) were chosen as an in vivo model to determine tumorigenesis in mammary glands where there is no functional ER $\alpha$  expressed. Incubation of MCF-10A<sup>arom</sup> cells with androstenedione (10  $\mu$ M) for 24 h resulted in an increase in the levels of 4-hydroxyestrone as well as its detoxicated product, 4-OHE-NAcCys, which was blocked by letrozole. These results indicate that breast epithelial cells have the enzymes that are required to convert estrogen to the genotoxic products. Overexpression of aromatase enhances in situ production of estrogen and its genotoxic metabolites. The ERKO/Wnt-1 mice were ovariectomized at 15 days. E<sub>2</sub> was administered through s.c. silastic capsules. Predicted serum concentrations of E<sub>2</sub> delivered by silastic implants were 80 and 240 pg/ml. By 11 months, tumor incidences were 50% in animals receiving high dose E<sub>2</sub> and 25% with lower dose. In the control group only 16% of animals developed tumors ( $p < 0.04$ ). By 18 months, 66% of animals receiving high dose E<sub>2</sub> developed mammary tumors. Our data provide the first direct evidence that estrogen may cause breast cancer through a non estrogen receptor mediated mechanism.

Submitted to the International Conference of Aromatase 2004, Edinburgh, UK